TEST PATIENT

TEST PHYSICIAN



P: 1300 688 522

E: info@nutripath.com.au A: PO Box 442 Ashburton VIC 3142 GUa d'Y'HYgh'BUa Y

Sex ::

DR JOHN DOE

111 CLINIC STF 99H

DUHY Collected: 00-00-0000 7@B=7 GI 6I F 6 J=7 " \$\$\$
111 H9GH ROAD TEST SUBURB

@AB =8: **00000000** UR#:0000000

ENDOCRINOLOGY SALIVA								
SALIVA	Result	Range	Units					
Male Hormone Profile-Extensive								
DHEAS.	18.9	5.0 - 30.0	nmol/L					
Androstenedione	1.4	1.0 - 10.7	nmol/L					
Testosterone.	389.0	100.0 - 720.0	pmol/L					
Estradiol (E2)	10.0		pmol/L					
Estrone (E1)	18.0	9.6 - 20.0	pg/mL					
Androstenedione/E1 Ratio	0.08	0.05 - 1.10	RATIO					

DR JOHN DOE



GUa d'Y HYgh BUa Y

@AB =8: 00000000 UR#:0000000

Sex::

111 CLINIC STF 99H DUNY Collected: 00-00-0000

7@=B=7 GI 6I F6 J=7 " \$\$\$ 111 H9GH'ROAD'TEST SUBURB

P: 1300 688 522 E: info@nutripath.com.au A: PO Box 442 Ashburton VIC 3142

Saliva Hormone Comments

SALIVARY HORMONE REFERENCE RANGES: (NOT ON HRT - BASELINE)

1	ı	E2	I	E1	I	E3	I	Progesterone	I	DHEAS
FEMALE Follicular Mid-Cycle Luteal	 	<18 11-29 <18	 	9.6-20 9.6-20 9.6-20	 	15-29 15-29 15-29	 	<318 - 318-1590		
Post Men. <6 9.6-20 1-41 <159 Premenopausal, no oral contraceptives Premenopausal, with oral contraceptives									 	<6.5 2.5-25.0 2.0-8.0
MALE		<6	1	9.6-20	1	16-25		<159	1	5.0-30.0

TARGET	ARGET REFERENCE RANGES:				(ON HRT - 24hr post last dose)						
		E2	 	E1	 	Е3	1	Progesterone	-	Testosterone Age Dpndt	1
Oral Patch Cream/	 Gel	7-73 4-18 37-184	 	- - -	 	69-139 - 1040-1734	 	318-1590 - 3180-31797	•	277-867 347-1734	1

SALIVA ESTRONE (E1) is produced primarily from androstenedione originating from the gonads or the adrenal cortex. In premenopausal women, more than 50% of the E1 is secreted by the ovaries. In prepubertal children, men and non-supplemented postmenopausal women, the major portion of E1 is derived from peripheral tissue conversion of androstenedione. Interconversion of E1 and E2 also occurs in peripheral tissue. Bioassay data indicate that the estrogenic action is much less than E2. E1 is a primary estrogenic component of several pharmaceutical preparations, including those containing conjugated and esterified estrogens. In premenopausal women E1 levels generally parallel those of E2. After menopause E1 levels increase, possibly due to increased conversion of androstenedione to E1.

SALIVA E1 level is within range for a male, indicating minimal aromatisation of androgens to E1.

ELEVATED E2 LEVEL:

Saliva E2 level is elevated for a male and suggestive of aromatisation of androgens to estrogens. Suggest using 5% transdermal Chrysin and / or 50mg Zinc.

SALIVA DHEAs level is adequate and within range. SALIVA FREE TESTOSTERONE level is adequate and within range.

Tests ordered: 1008